

Phytochemicals and Biological Activities of *Flueggea virosa* (Phyllanthaceae) Used in the Traditional Treatment of Benign Prostatic Hyperplasia in Mali

Adama Dénou^{1,2,*}, Mahamane Haïdara¹, Flacoro Diakité¹, Sékou Doumbia¹,
Daouda Lassine Dembélé¹, Rokia Sanogo^{1,3}

¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, University of Sciences, Techniques and Technologies of Bamako (USTTB), Bamako, Mali

²Department of Pharmacognosy and Traditional Medicine, University of Jos, Jos, Nigeria

³Department of Traditional Medicine, National Institute in Public Health, Bamako, Mali

Email address:

denouadamab@gmail.com (A. Dénou), smydos@yahoo.fr (M. Haïdara), diakiteflacoro@yahoo.fr (F. Diakité), s2doumbia@yahoo.fr (S. Doumbia), drdembeled@gmail.com (D. L. Dembélé), rosanogo@yahoo.fr (R. Sanogo)

*Corresponding author

To cite this article:

Adama Dénou, Mahamane Haïdara, Flacoro Diakité, Sékou Doumbia, Daouda Lassine Dembélé, Rokia Sanogo. Phytochemicals and Biological Activities of *Flueggea virosa* (Phyllanthaceae) Used in the Traditional Treatment of Benign Prostatic Hyperplasia in Mali. *Journal of Diseases and Medicinal Plants*. Vol. 7, No. 4, 2021, pp. 119-126. doi: 10.11648/j.jdmp.20210704.14

Received: September 24, 2021; **Accepted:** October 21, 2021; **Published:** November 29, 2021

Abstract: Worldwide, benign prostatic hyperplasia (BPH) is common in men over the age of 50. Hence there is a phytotherapy for BPH. In Mali, many medicinal plants are used in the traditional treatment of BPH. The current work aimed to characterize secondary metabolites and anti-radical constituents, to determine the acute toxicity and to evaluate the analgesic and anti-inflammatory activities of aqueous extracts from the leaves and stems of *Flueggea virosa*. Colour and precipitation reactions in tubes were performed to characterize the chemical constituents of the extracts. Anti-radical substances were determined through the free radical reduction method of 1,1 diphenyl, 2-picryl hydrazyl radical. The acute toxicity was performed *in vivo* using female mice. *In vivo* analgesic activity was determined through inhibition of induced pain in mice using 0.6% acetic acid. *In vivo* anti-inflammatory activity was determined by inhibiting inflammation caused with carrageenan in mice. The main chemical constituents of aqueous extracts were flavonoids, tannins and saponins. Polar extracts contained anti-radical substances. These extracts exhibited analgesic and anti-inflammatory activities. The decoction extracts administered orally in mice at the dose of 2000 mg/kg did not provoke toxic effects and mortality. The stem extract at 100 mg / kg exhibited the best analgesic activity with 64.82% inhibition and anti-inflammatory activity with 59.07% inhibition. These first results justify the traditional use of the plant in the medical management of BPH and will help to make data for the development of a new improved traditional medicine (ITM) based on the stems of *Flueggea virosa*.

Keywords: *Flueggea virosa*, Analgesic and Anti-inflammatory, Anti-radical Substances, Benign Prostatic Hyperplasia, Mali

1. Introduction

Benign prostatic hyperplasia (BPH) is a common disease in older men, with lower urinary tract symptoms caused by hyperplasia of the prostatic epithelium and stromal cells [1, 2]. Essential androgens for prostate growth during early development, puberty, and aging have been documented to play a role in enlarged prostate associated with other factors such as

aging, diet, increased inflammatory mediators, hormones and oxidative stress [2]. Benign prostatic hyperplasia (BPH) is a histologic concept, sometimes with symptoms in the lower urinary tract: frequent, imperious and incomplete urination, weakness of the jet, dysuria and nocturia [3]. Currently, the main therapeutic drugs for BPH are 5-alpha reductase inhibitors and beta blockers [4, 5], such as finasteride and terazosin. However, these drugs have side effects such as fatigue, hypotension, ejaculation disorders, dysfunction and an increased risk of

prostate fibrosis [5, 6]. Furthermore, surgical treatment is another powerful remedy for BPH [7].

Nowadays, this option poses a lot of complications and the possibility of recurrence. Thus, the number of patients undergoing surgery for BPH is gradually decreasing [8]. Therefore, finding a curative drug that can effectively treat BPH with few side effects is critical. Herbal agents have gained considerable attention for the treatment of BPH and prostate cancer because they are thought to be safer, cost effective and have fewer side effects than conventional alternatives, especially in developing countries [9, 10]. New theories support that many potential herbs may inhibit the action of an essential hormone-regulating enzyme that converts testosterone to dihydrotestosterone, a process considered important in the development of an enlarged prostate and prostate cancer [11].

The use of complementary and alternative medicine in the management and treatment of BPH has shown hope and is gaining popularity [12]. Some drugs like saw palmetto, *Cucurbita pepo* (pumpkin) have been shown to suppress the level of prostate specific antigen (PSA) in the blood [11]. In addition, studies of randomized trials and meta-analyses through recent literature suggest clinical efficacy and good tolerance of certain preparations, in particular extracts of *Serenoa repens* and also of *Pygeum africanum*, produced with high concentrations of β -sitosterol, and pumpkin seeds [13]. A recent literature review undertaken in Nigeria identified African medicinal plants or not with anti-BPH activity in animals [12]. These plant species belonging to different families were *Celosia argentea* F. Cristata (L) (Amaranthaceae), *Citrullus lanatus* (Thumb) (Cucurbitaceae), *Croton membranaceus* (Euphorbiaceae), *Cucurbita pepo* L. (Cucurbitaceae), *Raphia hookeri* G. Mann & H. A. Wendl. (Arecaceae / Palmaceae), *Secamone afzelii* (Asclepiadaceae), *Solanum macrocarpon* L. (Solanaceae), *Telfairia occidentalis* (Hook. F) (Cucurbitaceae), *Trichosanthes cucumerina*.var. *anguina* (Cucurbitaceae) and *Zapoteca portoricensis* (Jacq.) H. M (Fabaceae). These plants contain bioactive constituents and antioxidant agents such as flavonoids, terpenoids, saponins and steroids which contribute to their anti-inflammatory properties and thus to their anti-BPH activities [12]. Overall, the use of herbal remedies for benign prostatic hyperplasia is aimed at reducing symptoms associated with this disease. In Mali many medicinal species are used in the traditional treatment of BPH. The chemistry and biological activities of some of these plants including *Acanthospermum hispidum*, *Curculigo pilosa*, *Piliostigma thonningii* and *Spilanthes uliginosa* have been documented [14, 15].

Flueggea virosa (Roxb. ex Willd.) Royle belonging to the Phyllanthaceae family is a shrub two to three meters long, with many branches starting from the base, thus forming an ellipse. It is used in traditional medicine in Mali and elsewhere to manage several diseases including diarrhoea, rheumatism, malaria, liver disease, inflammation and pain. The extracts of this plant also find their use in certain disorders of urinary and genital tracts [16]. The literature shows that this plant is chemically rich in alkaloids, triterpenoids, tannins, flavonoids, resins, steroids, cardiac

glycosides, and anthraquinones [17]. The presence of these constituents confers to this plant a variety of pharmacological activities such as analgesic, anti-inflammatory, aphrodisiac, sedative, anti-arrhythmic, antidiabetic, antimalarial, anti-HIV, anti-hepatitis C, anti-diarrheal, cytotoxic, anti-microbial, antifungal, antioxidant, and laxative properties [17].

With the view for developing a phytomedicine for the management of BPH in Mali, the current work aimed to evaluate the analgesic, anti-inflammatory and antioxidant activities of aqueous extracts from the leaves and stems of *Flueggea virosa* used in the traditional treatment of BPH in Mali.

2. Materials and Methods

2.1. Plant Material

The stems and leaves of *Flueggea virosa* were collected in December 2012 by Mr. Oumar Diakit  in Fouladougou-bangassi in the Kita circle of Mali.

The samples arrived in Bamako the same day and were identified at the Department of Traditional Medicine (DMT) of the National Institute for Public Health Research (INRSP) then dried for a week under shade at room temperature in the drying room of the DMT. After drying the two samples were pulverized using a Forplex type F1 mill available at DMT. The powders obtained were used for extractions, and phytochemical and biological investigations.

2.2. Animals

For this work authors used white mice male and female weighing between 18 and 35 g, of no consanguineous strain from CF1 (Carworth Farms strain 1) line of OF1 (Oncins France strain 1) race supplied by the animal facility of the National Support Center for Disease Control (CNAM) in Bamako. The animals were kept under standard environmental conditions. They had a 12 hour light /dark cycle. Animals were fed using mouse food made at DMT with free access to tap water.

2.3. Ethical Statement

The experimental procedures and techniques used in the study were in accordance with accepted principles for laboratory animal use and care. All protocols and techniques used in this study were approved by the National Institute for Public Health Research Ethic Committee.

2.4. Extractions

Different types of extracts have been made for antioxidant activity and biological tests.

2.4.1. Aqueous Extracts

10% Extemporaneous Decoction

Authors mixed 10 g of the drug and 100 ml of distilled water together in a flask then put the flask on a warmer and boil for 15 minutes. After cooling, the extract was filtered through a 40x40 compress. The volume of the filtrate was

measured and then used directly for the tests.

10% Decoction

Authors mixed 100 g of the drug in 1000 ml of distilled water and boiled for 15 minutes. After cooling, the extract was filtered through a compress and squeezed. Authors obtained a final volume of 690 ml for the leaves and 700 ml for the stems. These final volumes were concentrated in a rotary evaporator and then frozen before being lyophilized in a freeze dryer. The yield was calculated for each extraction.

10% Infusion

Authors mixed 100 g of the drug in 1000 ml of boiling distilled water. After cooling we filtered on a compress and squeezed. Authors obtained a final volume of 830 ml for the leaves and 820 ml for the stems. These volumes were concentrated and then frozen before being lyophilized. The yields were then calculated.

2.4.2. Hydroethanolic (70% EtOH and 30% EtOH)

Extractions

Authors mixed 5 g of drug and 50 ml of solvent in a flask containing a magnetic rod then put this flask on the stirrer for 1 hour. Authors then filtered, evaporated to dryness in a rotary evaporator, weighed the extracts, then their yields were calculated.

2.5. Phytochemical Screening

Certain major groups of secondary metabolites have been investigated in drugs according to conventional methods of characterization. Tannins and polyphenols were identified by the FeCl₃ test; flavonoids by reaction to cyanidin; saponins by the foam test; the triterpenes and steroids by the Liebermann-Burchard test and finally the alkaloids by the tests of Mayer and Dragendorf [18, 19].

For the characterization of the anti-free radical constituents, the plates were revealed with the methanolic solution of 1,1 Diphenyl-2-Picryl-Hydrazyl (DPPH), in the proportion 2 mg/10 ml. When present the anti-radical constituents appear as yellow spots on a purple background.

2.6. Biological Activities

2.6.1. Determination of Acute Toxicity

The acute oral toxicity test was performed according to the guidelines of the Organization for Economic Co-operation and Development (OECD) [20]. The mice were fasted for 18 hours with free access to water. With a weight varying between 19 and 25 g, they were divided at random into five groups of three female mice and treated orally with a single dose of dry aqueous extract from the leaves or stems of *F. virosa* (2000 mg/kg), 20% extemporaneous extract of each plant organ (20 ml/kg) or distilled water (20 ml / kg) used as negative control. After administration, the animals were observed for the first 2 hours before feeding them and then observed for another two hours to record cases of immediate deaths or cases of behavioural change relating to the skin, hair, eyes, mucous membranes, respiration etc and once a day on 14 days to record late deaths.

2.6.2. Analgesic Activity

Analgesic activity was tested using 30 male mice (weighing 22-34 g). The animals were randomized into six groups of five mice. Groups I and II were treated orally with distilled water (20 ml/kg) used as negative control group and Paracetamol (100 mg/kg) used as reference drug respectively. Groups III and IV received orally 20 ml/kg of 10% extemporaneous decoction from the leaves and stems of the plant respectively. While groups V and VI were respectively treated orally with 250 mg/kg of dry 10% decoction extract from leaves and with 100 mg/kg of dry 10% decoction extract from stems. One hour after these treatments, the animals received by intra-peritoneal (IP) injection 0.6% acetic acid solution at a dose of 10 µl/g of body weight. Immediately after their injection, the mice were placed in plastic cages, observed to count the twists performed by each mouse during the 25 minutes following the intra-peritoneal injection of acetic acid as described by Siegmund et al. [21] with a few modifications. The time to onset of the first twist and the number of twists were noted for each mouse.

The mean value for each group was calculated and compared to the negative control group. The percentages of inhibition were calculated using the following formula:

$$\% \text{ Inhibition} = \frac{(W_c - W_t) \times 100}{W_c} \quad (1)$$

W_c represents the mean of the number of contortions of the mice in the negative control group;

W_t is the mean of the number of contortions of the mice in the groups treated with extracts and Paracetamol.

2.6.3. Anti-inflammatory Activity

The acute inflammation was induced by the injection of carrageenan under the plantar fascia of the right hind paw of mice as described by Winter et al [22] with slight modifications. Male mice were randomized into six groups of five animals. Groups I and II were treated orally with distilled water (20 ml/kg) used as negative control group and indomethacin (10 mg/kg) used as reference drug respectively. Groups III and IV received orally 20 ml/kg of 10% extemporaneous decoction from the leaves and stems of the plant respectively. While groups V and VI were respectively treated orally with 250 mg/kg of dry 10% leaf decoction and 100 mg/kg of dry 10% stem decoction. One hour after the treatments 0.025 ml of a 1% carrageenan suspension in physiological saline was injected in the right hind paw of each animal to induce the œdema. The paw thickness was measured using a digital calliper. The measurements were determined at 0 h or initial thickness (E₀) before injection of the carrageenan and after the injection of the carrageenan, the thickness of the paw was then measured every hour until the fifth hour (E_T). The difference between E_T (1, 2, 3, 4 and 5 h) and E₀ was considered as the œdema value. The average paw thickness for each group was calculated and compared to the negative control group according to the following formula.

$$\% \text{ œdema} = \frac{(E_T - E_0) \times 100}{E_0} \quad (2)$$

E0=Initial thickness of the mouse paw; ET=Thickness of the paw after injection of carrageenan and treatment

The percentages of inhibition were then calculated using the following formula:

$$\% \text{ Inhibition} = \frac{(P_0 - PT) \times 100}{P_0} \quad (3)$$

P0=% increase of the paw in the negative control; PT=% increase of the paw in the treated batch.

2.7. Statistical Analysis

The results were expressed as Mean \pm SD. The data were analysed by Student's t test. A value of P<0.05 was considered significant.

3. Results

3.1. Yields of Aqueous and Hydroalcoholic Extracts

The yields of the various extractions with polar solvents on the leaves and the stems are reported in Table 1. The yields of the leaf extracts were higher than those from the stem. The best yield was 41.60% for the 30% ethanolic extract from the leaves.

3.2. Phytochemical Data

The information on the chemical groups sought in the two parts analysed is noted in Table 2. The anti-free radical

substances present in the polar extracts from the two organs of the plant are illustrated by yellow spots in Figure 1.

Table 1. Yields of different extractions with polar solvents on the leaves and stems of *Flueggea virosa*.

Extracts	Yields (%)	
	Leaf	Stem
10% Decoction	25.35	8.56
10% Infusion	25.14	10.70
10% Extemporaneous Decoction	28.60	8.70
70% EtOH	22.60	14.40
30% EtOH	41.60	16.00

Table 2. Secondary metabolites analysed by tube reactions.

Chemical Groups	Results	
	Leaf	Stem
Flavonoids	++	+++
Alcaloids (Mayer-Dragendorff)	-	-
Tannins (reaction with FeCl ₃)	++++	+++
Gallic tannins (reaction of Stiasny)	++	+++
Catechic tannins	-	+++
Steroids and triterpenes	++	+++
Anthocyanins	-	-
Leucoanthocyanins	+	+
Saponins	±	±

Key: ++++=Highly abundant; +++=Abundant; ++=Moderate; +=Low; ±=Trace; -=Absent.




Figure 1. Chromatogram showing the anti-free radical constituents stained in yellow on the plate revealed with DPPH after migration in Butanol: Acetic acid: Water (60:15:25).

Key: 1=10% leaf decoction, 2=10% leaf infusion, 3=10% leaf Extemporaneous decoction, 4=70% leaf ethanol extract, 5=30% leaf ethanol extract, 6=10% stem decoction, 7=10% stem infusion, 8=10% stem Extemporaneous decoction, 9=70% stem ethanol extract and 10=30% stem ethanol extract.

3.3. Biological Activities

3.3.1. Acute Toxicity Data

From our experimental conditions, the 20% extemporaneous decoction (20 ml/kg) and the dry aqueous extracts at a dose of 2000 mg / kg of the leaves or stems of *Flueggea virosa* showed no signs of toxicity or mortality after 14 days.

3.3.2. Analgesic Activity

The results of the analgesic activity of the 10% extemporaneous and dry aqueous extracts of the leaves and stems of *Flueggea virosa* on the pain caused by acetic acid are reported in Table 3. Overall, all the extracts tested showed the analgesic activity. However, it was the dry extract of the stems that led to the best activity with 64.82% inhibition of pain.

Table 3. Analgesic activity of extemporaneous and dry aqueous extracts of the leaves and stems of *Flueggea virosa* on pain caused by acetic acid.

Treatments (doses/kg)	Number of contorsions (M ± SD)	% Inhibition
Distilled Water (20 ml/kg)	61.40 ± 01.14	----
10% Leaf Extemporaneous Decoction (20 ml/kg)	37.00±05.61**	39.73
Dry Leaf Decoction Extract (250 mg/kg)	18.80±08,10**	69.38
10% Stem Extemporaneous Decoction (20 ml/kg)	30.60±09.23**	50.16
Dry Stem Decoction Extract (100 mg/kg)	21.60±06.02**	64.82
Paracetamol (100 mg/kg)	13.00±05.91**	82.98

M=Mean from six mice; SD=Standard Deviation, t-Student *significant when $p<0.05$; **Highly significant with $p<0.01$ compared to negative control group treated with distilled water

3.3.3. Anti-inflammatory Activity

The data on the anti-inflammatory effects of the aqueous extracts (extemporaneous and dry) of the leaves and stems of *F. virosa* are shown in Tables 4-7 below.

Effects of extemporaneous extracts

Carrageenan caused significant inflammation to the paw of the animals (Table 4). The extracts acted better on inflammation in related time with better activity observed at the fifth hour by the extract from the stems with 54.57% inhibition of inflammation (Table 5).

Table 4. Effects of extemporaneous extracts on oedema induced by injection of carrageenan in mice.

Treatments (doses/kg)	Swelling Degree of paw (M ± SD) in mn over the time					
	E0	E1	E2	E3	E4	E5
Distilled Water (20 ml/kg)	0.76±0.2	1.95±0.2	2.08±0.3	2.03±0.2	2.18±0.2	2.15±0.5
Leaf Extract (20 ml/kg)	0.68±0.2	1.96±0.3	1.66±0.2	2.05±0.3	2.06±0.3	1.80±0.4
Stem Extract (20 ml/kg)	1.39±0.2	1.85±0.3	1.89±0.3	1.76±0.2	1.96±0.3	1.89±0.3

M=Mean from 5 mice; DS=Standard Deviation; E=Edema over the time

Table 5. Percentage inhibition of oedema by extemporaneous extracts according to negative control.

Treatments (doses/kg)	% Inhibition of Oedema					
	0h	1h	2h	3h	4h	5h
Distilled Water (20 ml/kg)	-	-	-	-	-	-
Leaf Extract (20 ml/kg)	-	AI	17.15*	23.22*	34.60*	48.52**
Stem Extract (20 ml/kg)	-	14.65*	34.30*	40.71**	47.28**	54.57**

E=Edema over the time; *Significant when $p<0.05$; **Highly significant when $p<0.01$ compared to negative control group treated with distilled water. AI=Absence of inhibition.

Effects of dry aqueous extracts

The activity was organ and dose dependent according to the swelling degree (Table 6). From the third hour we saw

good anti-inflammatory activity with the dry stem extract which was 59.07% inhibition even exceeding that of indomethacin used at 10 mg / kg (Table 7).

Table 6. Effects of dry extracts on oedema induced by injection of carrageenan in mice.

Treatments (doses/kg)	Swelling Degree of paw (M ± SD) in mm over the time			
	E0	E1	E2	E3
Distilled Water (20 ml/kg)	1.34±0.1	2.27±0.3	2.6±0.5	2.31±0.4
Leaf Extract (250 mg/kg)	1.33±0.1	2.13±0.2	2.64±0.1	2.29±0.2
Stem Extract (100 mg/kg)	1.24±0.1	2.11±0.2	2.23±0.4	2.09±0.2

M=Mean from 5 mice; DS=Standard Deviation; E=Edema over the time

Table 7. Percentage inhibition of oedema by the dry extracts and the reference drug according to the negative control.

Treatments (doses/kg)	% Inhibition of Oedema			
	0h	1h	2h	3h
Distilled Water (20 ml/kg)	-	-	-	-
Leaf Extract (250 mg/kg)	-	AI	AI	12.04*
Stem Extract (100 mg/kg)	-	32.40*	54.60**	59.07**
Indomethacin (10mg/kg)	-	6.14*	24.40*	26.31*

*Significant with $p<0.05$; **Highly significant with $p < 0.01$ compared to negative control group treated with distilled Water. AI=Absence of Inhibition.

4. Discussion

Flueggea virosa is a plant whose roots are used in traditional medicine in men with benign prostatic hyperplasia, but the stems and leaves were chosen in order to keep the plant alive.

The leaf extracts were greenish in colour and sticky in consistency while the stem extracts were yellowish and less sticky. The yields obtained for the leaves were much higher than those for the stems. This would probably be related to a rapid release of the polar constituents from the leaves when they are in contact with the polar solvents.

The leaves and stems were rich in secondary metabolites such as flavonoids, tannins, sterols and triterpenes, leucoanthocyanins but with traces of saponins. Tube reactions do not revealed the presence of alkaloids in the two samples.

The leaves were rich in coumarins, carotenoids and mucilages which on the other hand, were absent in the stems [23]. The colour and precipitation reactions used in this present work failed to demonstrate alkaloids in the investigated samples. However, they have been demonstrated in leaf extracts by thin layer chromatography using Dragendorff's reagent [23]. This result goes in the same direction like the one obtained by recent work carried out at DMT on leafy stems of *Flueggea virosa* [24]. In Nigeria the presence of alkaloids, tannins and flavonoids in the organs of *Flueggea virosa* has been revealed in methanolic extracts from the roots and leaves of *Flueggea virosa* [25, 26]. Also in Nigeria, scientists have found in addition alkaloids, saponins, flavonoids, anthraquinones and steroids in the leaf methanolic extract of the plant [27].

Extracts from the leaves and stems showed the presence of anti-free radicals by scavenging the DPPH radical. However, these anti-free radicals were more pronounced with the hydroalcoholic extracts of the leaves. The presence of anti-free radical substances could be explained by the richness of the organs analysed in polyphenolic compounds. Numerous studies have shown the anti-free radical activity of extracts or isolated molecules of *Flueggea virosa* [28-30].

Based on the current experimental conditions, the 20% extemporaneous decoction extracts and the dry 10% aqueous extracts at 2000 mg / kg showed no signs of toxicity after 14 days. Many studies have shown the low toxicity of *Flueggea virosa*. In 2008 Magaji and his team found some LD₅₀ of 774.6 mg / kg and greater than 5000 mg / kg after respective administration of root methanolic extracts intraperitoneally and orally in mice [25]. Other previous authors found some LD₅₀ of 1265 mg / kg and 1264.9 mg / kg, respectively, after intraperitoneally administration of leaf methanolic extract, respectively in mice and rats [26, 27]. The toxicity mentioned in Kheraro (1974) and reported by Paris *et al.*, 1955 at doses of 2.5 to 5 g / kg in mice could depend on the source of their sample [16]. In Mali Traoré *et al.* found that the toxicity dose of leafy stems of *F. virosa* was greater than 2000 mg / kg [24].

Regarding analgesic and anti-inflammatory activities, we found that most of the extracts exhibited analgesic and anti-inflammatory activities, but the best activities were those of the dry stem aqueous extract at 100 mg / kg per oral route with 64.82% as the percentage inhibition of pain and 59.07% against inflammation. These effects could be due to the presence of saponins. Indeed, according to Catier and Roux, 2007, some saponins are endowed with analgesic and anti-inflammatory properties [31].

The current results confirmed those of Magaji and coworkers who showed that the root methanolic extract (6.25; 12.5 and 25 mg / kg) administered intraperitoneally inhibited the acetic acid-induced pain in a dose-dependent manner but the strong activity was exhibited by the dose of 25 mg / kg, alleviating the pain induced by formalin. It increased significantly (P <0.01) the latency time for hot plate-induced pain and inhibited carrageenan-induced inflammation (P <0.05) in mice [25]. In Nigeria, researchers found similar results with leaf methanolic extract (25, 50 and 100 mg / kg) administered intraperitoneally in mice [26]. In 2012 Ezeonwumelu and his colleagues found analgesic activity with root extracts at doses of 200 and 400 mg / kg administered orally in rats [32]. Recent work carried out in Mali has shown that the leafy branches of *F. virosa* exhibited analgesic effects and remarkable anti-inflammatory activities at 100 and 200 mg / kg [24].

The major goal of all drugs used against BPH is to reduce the number of symptoms associated with this condition in order to improve the patient's quality of life. The imbalance between oxidative stress and antioxidants plays an important role in the development of benign prostatic hyperplasia [33]. Although the mechanism by which oxidative stress causes inflammation remains unclear, it is clear that oxidative stress is associated with many chronic inflammatory diseases. A general consensus shows that polyphenols are powerful anti-inflammatory and antioxidant factors through complex mechanisms. *In vitro* studies on BPH cell lines as well as several *in vivo* works using animal models have shown antioxidant and anti-inflammatory effects [34]. Furthermore, all pain has an inflammatory profile made up of inflammation mediators present in this pain syndrome [35]. The late diffuse pain induced in mice by intraperitoneally administration of acetic acid was manifested by stretching movement of the hind paw and torsions of the dorso-abdominal musculature. Prior studies on medicinal plants from two different countries found that many African people used medicinal plants to treat pain [36, 37]. In addition a pharmacological investigation confirmed the analgesic activity against acetic acid induced-pain in mice of aqueous extracts from four medicinal plants used in Mali and Togo [38]. Elsewhere the extracts of *F. virosa* tested on this pain significantly reduced its symptoms. Authors have reported that the reduction in pain induced by acetic acid in mice may be due to the presence of polyphenols such as flavonoids [39]. Thus present extracts, all rich in polyphenols with antioxidant, anti-inflammatory and analgesic properties,

could act on the functional component of the prostate by reducing its volume and the pressure exerted by its muscles on the urethra. This could explain how extracts of *Flueggea virosa* may have a beneficial effect in the medical management of BPH.

5. Conclusion

This work aimed to validate and justify the traditional use of the leaves and stems of *Flueggea virosa* in the management of benign prostatic hyperplasia. Indeed, these samples contain polyphenolic compounds (flavonoids, tannins, etc.) with anti-free radical properties.

The dry extract of the stems (100 mg / kg) demonstrated the best analgesic and anti-inflammatory activities. The results of acute toxicity support the safety of the aqueous (extemporaneous and dry) extracts used at the therapeutic dose in the management of benign prostatic hyperplasia. These effects may therefore be beneficial in the medical management of BPH. This work will contribute to the development of a new improved traditional medicine (ITM) based on the stems of *Flueggea virosa*.

Conflict of Interest

All the authors do not have any possible conflicts of interest.

References

- [1] Saito M, Tsounapi P, Oikawa R, Shimizu S, Honda M, Sejima T, Kinoshita Y, Tomita S (2014). Prostatic ischemia induces ventral prostatic hyperplasia in the SHR; possible mechanism of development of BPH. *Scientific reports*. 4 (1): 1-7.
- [2] Bostanci Y, Kazzazi A, Momtahan S, Laze J, Djavan B (2013). Correlation between benign prostatic hyperplasia and inflammation. *Current Opinion in Urology*. 23: 5-10.
- [3] Anonymous. Medical management of benign prostatic hyperplasia. *Therapeutics Letter* may/june 1997; 19a-b. <http://www.ti.ubc.ca/PDF/19.PDF>.
- [4] Sáez C, González-Baena AC, Japón MA, Giráldez J, Segura DI, Miranda G, Rodríguez-Vallejo JM, González-Esteban J, Torrubia F (1998). Regressive changes in finasteride-treated human hyperplastic prostates correlate with an upregulation of TGF- β receptor expression. *The Prostate*. 37 (2): 84-90.
- [5] Tract Symptoms/Benign Prostatic Hyperplasia Guidelines Committee (2017). Singapore urological association clinical guidelines for male lower urinary tract symptoms/benign prostatic hyperplasia. *Singapore medical journal*. 58 (8): 473-480.
- [6] Kim HK, Zhao C, Choi BR, Chae HJ, Kim DS, Park JK (2013). Is transforming growth factor- β signaling activated in human hypertrophied prostate treated by 5-alpha reductase inhibitor?. *Disease markers*. 35 (6): 679-85.
- [7] Adewale AO, Adedeji SI, Olusola OO, Sulaiman SO, Victoria IO (2021). Ethanolic Extract of Combined *Cynodon dactylon* and *Mimosa pudica* Ameliorated Experimentally Induced Benign Prostatic Hyperplasia in Wistar Rats. *Journal of Infertility and Reproductive Biology*. 9 (1): 22-26.
- [8] Minutoli L, Rinaldi M, Marini H, Irrera N, Crea G, Lorenzini C, Puzzolo D, Valenti A, Pisani A, Adamo EB, Altavilla D (2016). Apoptotic pathways linked to endocrine system as potential therapeutic targets for benign prostatic hyperplasia. *International Journal of Molecular Sciences*. 17 (8): 1311.
- [9] Arora RP, Nayak RL, Malhotra V, Mohanty NK (2003). Role of herbal drugs in the management of benign prostatic hyperplasia: Clinical trial to evaluate the efficacy and safety of Himplasia. *Med Update*. 11: 55-58.
- [10] Caesar LK, Cech NB (2019). Synergy and antagonism in natural product extracts: when 1+1 does not equal 2. *Natural product reports*. 36 (6): 869-888.
- [11] Olamide AA, Olayemi OO, Demetrius OO, Olatoye OJ, Kehinde AA (2011). Effects of methanolic extract of *Citrullus lanatus* seed on experimentally induced prostatic hyperplasia. *European Journal of Medicinal Plants*. 4: 171-9.
- [12] Ikeyi AP, Ugwoke AO (2020). A Review of Medicinal Plants with Potential Anti-Benign Prostatic Hyperplasia (Bph) Activity Used In Nigeria. *IDOSR Journal of Biochemistry, Biotechnology and Allied Fields*. 5 (1): 64-71.
- [13] Dreikorn K (2002). The role of phytotherapy in treating lower urinary tract symptoms and benign prostatic hyperplasia. *World Journal of Urology*. 19: 426-435.
- [14] Diarra Y (2006). Etude de la phytochimie et des activités biologiques de *Acanthospermum hispidum* DC. (Asteraceae) et de *Curculigo pilosa* Schum et Thonn. (Hypoxidaceae) deux plantes utilisées dans le traditionnel de l'HBP. Thèse de pharmacie, FMPOS, Bamako, Mali.
- [15] Diabaté M, (2005). Etude de la phytochimie et des activités biologiques d'une recette traditionnelle composée de la partie aérienne de *Spilanthes uliginosa* et des fruits de *Piliostigma thonningii* utilisée dans le traitement de l'HBP au Mali. Thèse de pharmacie, FMPOS, Bamako, Mali.
- [16] Kerharo J, Adam JC (1974). Pharmacopée sénégalaise traditionnelle: Plantes médicinales et toxicologiques [Senegalese Traditional Pharmacopoeia: Medicinal and Toxicological Plants]. Paris Vigot et Frères Ed.
- [17] Renu, Narain S, Rawat A, Kaur J, Kumar S, Fatima N (2018). Taxonomy, phytochemistry, pharmacology and traditional uses of *Flueggea virosa* (Roxb. Ex Willd.) Royle: A Review. *International Journal of Life Sciences*. 6 (2): 579-585.
- [18] Bruneton J (2009). Pharmacognosie, phytochimie, plantes médicinales (4e éd.). Lavoisier.
- [19] Evans WC (2002). Pharmacognosy. Saunders Elsevier.
- [20] Organization for Economic Co-operation and Development (2001). OECD Guideline for the Testing of Chemicals: Acute Oral Toxicity – Acute Toxic Class Method. N°423.
- [21] Siegmund E, Cadmus R, Lu G (1957). A method for evaluating both non-narcotic and narcotic analgesics. *Proceedings of the Society for Experimental Biology and Medicine*. 95 (4), 729-731.
- [22] Winter CA, Risley EA, Nuss GW (1963). Anti-inflammatory and antipyretic activities of indo-methacin, 1-(p-chlorobenzoyl)-5-methoxy-2-methyl-indole-3-acetic acid. *Journal of pharmacology and Experimental Therapeutics*. 141 (3): 369-76.

- [23] Diakité F (2014). Etude de la chimie et des activités biologiques de *Flueggea virosa* (Roxb. Ex Willd.) Voigt (Euphorbiaceae) utilisée dans le traitement traditionnel de l'hypertrophie bénigne de la prostate. Thèse de pharmacie, FAPH, Bamako, Mali.
- [24] Traoré K, Haidara M, Dénou A, Kanadjigui F, Sogoba MN, Diarra B, Maiga S, Sanogo R (2019). Criblage Phytochimique et Activités Biologiques de Quatre Plantes Utilisées au Mali dans la Prise en Charge du Paludisme Chez les Enfants. *European Scientific Journal*. 15 (6): 210-226.
- [25] Magaji MG, Anuka JA, Abdu-Aguye I, Yaro A H, Hussaini IM (2008). Preliminary studies on anti-inflammatory and analgesic activities of *Securinega virosa* (Euphorbiaceae) in experimental animal models. *Journal of Medicinal Plants Research*. 2 (2): 1-7.
- [26] Yerima M, Magaji MG, Yaro AH, Tanko Y, Mohammed MM (2009). Analgesic and antiinflammatory activities of the methanolic leaves extract of *Securinega virosa* (Euphorbiaceae). *Nigerian Journal of Pharmaceutical Sciences*. 8 (1): 47-53.
- [27] Tanko Y, Okasha M A, Magaji MG, Yerima M, Yaro AH, Saleh MIA, Mohamed A (2008). Anti-diabetic properties of *Securinega virosa* (Euphorbiaceae) leaf extract. *African Journal of Biotechnology*. 7 (1): 022-024.
- [28] Sanogo R, Vassallo A, Malafronte N, Imperato S, Russo A, Piaz F (2009). New phenolic glycosides from *Securinega virosa* and their antioxydant activity. *Journal Natural Product Communications*. 4 (12): 1645–1650.
- [29] Sanogo B (2010). Etude des Activités antioxydantes et anticonvulsivantes de *Securinega virosa* et de *Pteleopsis suberosa* deux plantes médicinales utilisées dans le traitement traditionnel de l'HBP.
- [30] Chauke AM, Shai LJ, Mphahlele PM, Mogale MA (2012). Radical scavenging activity of selected medicinal plants from Limpopo province of South Africa. *African Journal of Traditional Complementary and Alternative Medicines*. 9 (3): 426-30.
- [31] Catier O, Roux D (2007). Cahier du préparateur en pharmacie. Botanique –Pharmacognosie – Phytothérapie. 3ème Edition. Collection Porphyre.
- [32] Ezeonwumelu JO, Omar AN, Ajayi AM, Okoruwa AG, Tanayen JK, Kiplagat DM, Okpanachi OA, Abba S, Ezekiel I, Onchweri AN, Okonkwo CO (2012). Phytochemical screening, acute toxicity, anti-inflammatory and antipyretic studies of aqueous extract of the root of *Flueggea virosa* (Roxb. ex Willd.) in rats. *International Journal of Pharmacy and Biomedical Sciences*. 3 (3): 128-35.
- [33] Minciullo PL, Antonino IA, Navarra M, Calapai G, Magno C, Gangemi S (2015). Oxidative Stress in Benign Prostatic Hyperplasia: A Systematic Review. *Urologia Internationalis*. 94: 249–254.
- [34] Mitsunari K, Miyata Y, Matsuo T, Mukae Y, Otsubo A, Harada J, Kondo T, Matsuda T, Ohba K and Sakai H (2021). Pharmacological Effects and Potential Clinical Usefulness of Polyphenols in Benign Prostatic Hyperplasia. *Molecules*. 26 (450): 1-20.
- [35] Omoigui S (2007). The Biochemical Origin of Pain: The origin of all Pain is Inflammation and the Inflammatory Response. PART 2 of 3 – Inflammatory Profile of Pain Syndromes. *Med Hypotheses*. 69 (6): 1169–1178.
- [36] Denou A, Koudouvo K, Togola A, Aziati KY, Esseh J, Ajavon CA, Essien K, Aklikokou K, Sanogo R, Diallo D, Gbeassor M (2016). Traditional knowledge on antimalarial plants having analgesic properties, used in Togo Maritime Region. *The Journal of Ethnobiology and Traditional Medicine, Photon*. 126: 1160-70.
- [37] Dénou A, Koudouvo K, Togola A, Haïdara M, Dembélé SM, Ballo FN, Sanogo R, Diallo D, Gbeassor M (2017). Savoir traditionnel sur les plantes antipaludiques à propriétés analgésiques, utilisées dans le district de Bamako (Mali). *Journal of Applied Biosciences*. 112: 10985-95.
- [38] Denou A, Koudouvo K, Haidara M, Togola A, Sanogo R, Essien K, Aklikokou KA, Diallo D, Gbeassor M (2016). Activité analgésique de quatre plantes utilisées dans la prise en charge traditionnelle du paludisme au Mali et au Togo. *International Journal of Biological and Chemical Sciences*. 10 (3): 1342-9.
- [39] Lakache Z, Tigrine C, Aliboudhar H, Kamel A (2021). Composition chimique, activités anti-inflammatoire, antalgique et cytotoxique *in vivo* de l'extrait méthanolique des feuilles d'*Olea europaea*. *Phytothérapie*. 19 (2): 83-92.